


A detailed illustration of a fly, likely a housefly, resting on a large, green, veined leaf. The fly is shown from a side profile, with its wings spread slightly. The background is a solid, vibrant green.

## References

- doi:10.1016/j.cretres.2009.09.007 (<https://doi.org/10.1016/j.cretres.2009.09.007>).

- Cascoplecia**  
Temporal range:  
Late Aptian to Early Cenomanian
- |      |   |   |   |   |   |   |   |   |   |     |
|------|---|---|---|---|---|---|---|---|---|-----|
| PreЄ | Є | O | S | D | C | P | T | J | K | PgN |
|------|---|---|---|---|---|---|---|---|---|-----|



## Scientific classification

Kingdom:	Animalia
Phylum:	Arthropoda
Class:	Insecta
Order:	Diptera
Infraorder:	Bibionomorpha
Family:	† <b>Cascopleciidae</b> Poinar Jr., 2010
Genus:	† <b><i>Cascoplecia</i></b> Poinar Jr., 2010
Species:	† <i>C. insolitis</i>

**Binomial name**

†*Cascoplecia insolitis*  
Poinar Jr., 2010

---

Retrieved from "<https://en.wikipedia.org/w/index.php?title=Cascoplecia&oldid=959582363>"

---

**This page was last edited on 29 May 2020, at 14:00 (UTC).**

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.